

Knockdown of a metathoracic scent gland desaturase enhances the production of (E)-4-oxo-2-hexenal and suppresses female sexual attractiveness in the plant bug *Adelphocoris suturalis*

J. Luo*†, Z. Li*, C. Ma*, Z. Zhang‡, J. J. Hull§, C. Leit, S. Jin* and L. Chen*†

*National Key Laboratory of Crop Genetic Improvement and National Centre of Plant Gene Research, Huazhong Agricultural University, Wuhan, China; †Hubei Insect Resources Utilization and Sustainable Pest Management Key Laboratory, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China; ‡Hubei Key Laboratory of Quality Control of Characteristic Fruits and Vegetables, Hubei Engineering University, Xiaogan, China; and §U.S. Department of Agriculture, U.S. Arid Land Agricultural Research Center, Agricultural Research Service, Maricopa, AZ, USA

Abstract

Insect sex pheromones (SPs) are central to mate-finding behaviour, and play an essential role in the survival and reproduction of organisms. Understanding the roles, biosynthetic pathways and evolution of insect chemical communication systems has been an exciting challenge for biologists. Compared with Lepidoptera, little is known about the mechanisms underlying pheromone biosynthesis in Hemiptera. In this study, we isolated and characterized two new desaturase-like genes, termed *Asutdes1* and *Asutdes2*, from *Adelphocoris suturalis*, an important agricultural pest in China. Although the two genes encode an identical protein, Southern blot analysis revealed that they are duplicated genes. The *Asutdes2* transcript is more abundant than *Asutdes1* in the tissues tested, in particular the metathoracic scent gland and fat body. Silencing *Asutdes* expression in females by injecting double-stranded RNA

(*dsAsutdes*) against a portion of the coding sequence shared by the two genes enhanced the production of (E)-4-oxo-2-hexenal, a component of the *A. suturalis* SP blend, and dramatically suppressed the sexual attractiveness of *A. suturalis* females. We conclude that *dsAsutdes* is associated with the SP biosynthetic pathway in *A. suturalis*.

Keywords: *Adelphocoris suturalis*, desaturase, sex pheromones biosynthesis, (E)-4-oxo-2-hexenal, RNA interference (RNAi).

Introduction

Despite years of study, a complete understanding of the roles, biosynthetic pathways and evolution of insect chemical communication systems remains an ongoing challenge. In insects, sex pheromones (SPs) comprise a diverse group of chemical compounds that are central to mate-finding behaviour (Jurenka, 2004; Johansson & Jones, 2007). The chemical composition of the SP blends is highly specific even amongst closely related species (Symonds & Elgar, 2008), and in numerous species, changes in the SP profile that affect male attractiveness more broadly or differentially have been implicated as potential mechanisms underlying reproductive isolation and speciation (Phelan, 1992; Symonds & Elgar, 2008). Allelic variations in genes encoding SP biosynthetic enzymes that affect the composition of SPs and/or assortative mating have been well demonstrated in *Drosophila* (Coyne & Charlesworth, 1997; Chung *et al.*, 2014) and Lepidoptera (Roelofs *et al.*, 2002; Lassurance *et al.*, 2010; Buček *et al.*, 2015). However, how SP chemical diversity arises is a long-standing problem.

Studies in Lepidoptera, Diptera, Coleoptera and Hymenoptera that focused on elucidating the chemical composition of SPs have revealed a diversity of chemical structures (Symonds & Elgar, 2008). SPs are typically blends of alcohols, aldehydes, ketones, acetates

First published online 16 June 2017.

Correspondence: Lizhen Chen, National Key Laboratory of Crop Genetic Improvement and National Centre of Plant Gene Research, Huazhong Agricultural University, Wuhan 430070, China. Tel.: 027-87282130; e-mail: lzchen@mail.hzau.edu.cn

and long-chain hydrocarbons (10–30C), many of which contain carbon–carbon double bonds in varied positions and spatial orientations (El-Sayed, 2016). Although the chemical backbone of most pheromones arises from general biosynthetic pathways, the observed diversity is generated by a series of enzyme-based substrate modifications that include chain shortening and elongation, desaturation, reduction, oxidation, acetylation, aromatization, decarboxylation, and hydrolysis (Tillman *et al.*, 1999; Roelofs & Rooney, 2003). In moths, SPs are typically derived from fatty acid biosynthesis with pheromone biosynthetic pathway enzymes introducing various modifications including: fatty acid chain-shortening enzymes (Jurenka *et al.*, 1994; Tabata & Ishikawa, 2005) that use β -oxidation to shorten the pheromone precursors, fatty acid desaturases (Roelofs *et al.*, 2002; Moto *et al.*, 2004; Xue *et al.*, 2007; Wang *et al.*, 2010; Fujii *et al.*, 2011; Albre *et al.*, 2012; Groot *et al.*, 2014; Hagström *et al.*, 2014) that insert carbon–carbon double bonds into the fatty acyl chains, and fatty acid reductases (Moto *et al.*, 2003; Lassance *et al.*, 2013; Antony *et al.*, 2016) that convert fatty acyl precursors to their corresponding alcohols. The oxygenated functional groups, however, can also be further modified by oxidases and acetyltransferases (Knipple *et al.*, 2002; Jurenka, 2004). The molecular basis of lepidopteran pheromone biosynthesis has been effectively elucidated (Groot *et al.*, 2016), and numerous studies have been conducted with the goal of similarly understanding the pheromone biosynthetic pathways in Diptera (Chertemps *et al.*, 2006, 2007; González-Caballero *et al.*, 2014; Ng *et al.*, 2015) and Coleoptera (Sandstrom *et al.*, 2006; Rubi *et al.*, 2012; Keeling *et al.*, 2013; Song *et al.*, 2014). In contrast, despite numerous studies identifying hemipteran pheromone components and how different compounds affect insect behaviour, little is known about the underlying pheromone biosynthetic mechanisms.

Adelphocoris suturalis (Hemiptera: Miridae) is a polyphagous pest that was originally a secondary pest of cotton, but has since become a significant problem for cotton growing regions in China owing to a reduction in broad-spectrum insecticides following the widespread adoption of transgenic *Bacillus thuringiensis* (Bt) cotton (Lu *et al.*, 2008a; Li *et al.*, 2010). Recently, two female metathoracic scent gland (MTG)-derived SP components, (E)-4-oxo-2-hexenal (E4O2H) and hexyl hexanoate (HH) were identified in *A. suturalis* (Zhang *et al.*, 2014, 2016). Both compounds are synthesized and released by females around 7–10 days posteclosion (PE; Zhang *et al.*, 2011). The structure of these semiochemicals is similar to moth SPs (ie carbon–carbon double bonds and oxygenated functional groups comprising aldehydes and acetates). Thus, it can be hypothesized that pheromone biosynthesis in *A. suturalis* is catalysed by homologues of moth

pheromone biosynthetic enzymes. To identify potential *A. suturalis* SP biosynthesis-related genes, we mined previous *A. suturalis* MTG Illumina sequencing data (Luo *et al.*, 2014) for transcripts with high sequence similarity to key moth pheromone biosynthesis enzymes. We identified four candidate genes that showed higher expression in female *A. suturalis* MTGs during the calling period: *fatty acyl-coenzyme A (CoA) reductase*, *desaturase (des)*, *acetyltransferase* and *alcohol oxidase*. RNA interference (RNAi)-mediated silencing of *A. suturalis desaturase (Asutdes)* expression potentially suppressed the sexual attractiveness of females to males in the field; we thus hypothesized that *Asutdes* functions in *A. suturalis* SP biosynthesis. To test this hypothesis, Y-tube bioassays, field trapping experiments and gas chromatography mass spectrometry (GC-MS) analyses were performed to investigate the effects of RNAi-mediated gene silencing. We showed that suppressing *Asutdes* expression enhanced E4O2H production, changed the ratio of the *A. suturalis* SP blend, and clearly suppressed the sexual attractiveness of *A. suturalis* females to males.

Results

cDNA cloning and sequence analysis

Based on the *A. suturalis* transcriptome data (Luo *et al.*, 2014), we isolated two cDNAs encoding putative *Asutdes* sequences and designated them as *Asutdes1* and *Asutdes2*. Although the 5' untranslated region of the 1863-bp (*Asutdes1*) and 1780-bp (*Asutdes2*) transcripts differed in size and sequence composition (Fig. 1A), both genes contained the same 1077-bp open reading frame (ORF) encoding the same 358 amino acid residue protein. Southern blot analysis was performed to determine if the two *Asutdes* transcripts were derived from two independent genes or a single gene that underwent alternative splicing. The presence of two distinct bands on the blot indicated that *Asutdes1* and *Asutdes2* are two independent genes (Fig. 1B).

The deduced *Asutdes* amino acid sequence shares 61% sequence similarity with the *Helicoverpa zea* acyl-CoA Δ 9-desaturase (Fig. 2A) that catalyses formation of two minor but essential components of the *H. zea* pheromone blend (Rosenfield *et al.*, 2001). High similarity was also observed with a *Choristoneura rosaceana* metabolic Z9-desaturase (60% similarity) that acts on palmitic acid (Hao *et al.*, 2002), a *Trichoplusia ni* Δ 9-desaturase (58% similarity) with high specificity for stearic acid (Liu *et al.*, 1999), and the *Drosophila melanogaster* desaturase1 (58% similarity) responsible for the first desaturation step in fruit fly pheromone biosynthesis (Dallerac *et al.*, 2000). *Asutdes* also exhibited 50% sequence similarity with the bifunctional *Bombyx mori* Z11/10,12 fatty-acyl




Figure 1. Gene structure of *Adelphocoris suturalis* desaturase 1 (*Asutdes1*) and *Asutdes2*. (A) Schematic depiction of the gene structure for *Asutdes1* and *Asutdes2* with the regions selected for double-stranded RNA indicated. (B) A common region of the two *Asutdes* genes was labelled with digoxigenin and used as a probe for Southern blot analysis, in which *A. suturalis* genomic DNA was digested with *HindIII*-HF. The Southern blot analysis results showed that *Asutdes1* and *Asutdes2* are two independent genes. M, marker; ORF, open reading frame; RNAi, RNA interference; UTR, untranslated region. [Colour figure can be viewed at wileyonlinelibrary.com]

desaturase that functions in bombykol biosynthesis (Moto *et al.*, 2004) (Fig. 2A).

The predicted *Asutdes* protein shares several key features with other known insect desaturases, including three conserved histamine motifs (H1–H3), which are crucial for spanning the membrane bilayer and catalytic activity (Shanklin *et al.*, 1994; Haritos *et al.*, 2014), and four transmembrane regions (Fig. 2A). To assess the

phylogenetic relationship of *Asutdes* with other insect desaturases, a maximum-likelihood tree was generated using a representative set of desaturase sequences from 39 additional species. *Asutdes* sorted to a monophyletic clade with two plant bug desaturase sequences (Fig. 2B) away from the multiple desaturase subfamilies (ie $\Delta 5$, $\Delta 6$, $\Delta 9$, $\Delta 10$, $\Delta 11$, $\Delta 12$, $\Delta 14$) reported in moths, beetles and flies.




Figure 2. Sequence analysis of *Adelphocoris suturalis* desaturase (*Asutdes*). (A) Alignment of the predicted *Asutdes* protein and functional desaturase proteins from other insects. Black, grey and white backgrounds indicate amino acid conservation. The transmembrane domains are indicated by black lines above the sequence, and the three His domains are labelled H1–H3. Abbreviated species names correspond to: *Asut*, *A. suturalis* [Accession no. KX765194]; *Hzea*, *Helicoverpa zea* (Accession no. AF272343). *Tni*, *Trichoplusia ni* (Accession no. AAB92583); *Cros*, *Choristoneura rosaceana* (Accession no. AAN39697) *Bmor*, *Bombyx mori* (Accession no. BAD18122); *Dmel*, *Drosophila melanogaster* (Accession no. NP_731711); *Tcas*, *Tribolium castaneum* (Accession no. ADK13054). (B) Phylogeny of *Asutdes* with other insect desaturases. The unrooted maximum-likelihood tree was obtained using 86 sequences from 40 species. The confidence values at the edges are derived from 1000 rapid bootstrap replicates. Bootstrap values > 80% are shown on branches. Sequence abbreviations correspond to species names (see Table S1) with *Atha* (*Arabidopsis thaliana*, Accession no. NP_172098) used as the outgroup. The red triangle denotes *Asutdes*. Accession numbers can be found in Table S1. [Colour figure can be viewed at wileyonlinelibrary.com]




Figure 3. Tissue-dependent transcription pattern analysis of *Adelphocoris suturalis* desaturase 1 (*Asutdes1*) and *Asutdes2*. Relative expression levels of *Asutdes1* and *Asutdes2* transcripts in the head, metathoracic scent gland (MTG), midgut, ovary and fat body of females 10 days posteclosion. The values are expressed as the means \pm SEM based on three independent biological replicates. Different letters indicate significant differences ($P < 0.05$, one-way analysis of variance followed by Tukey's Honestly Significant Difference (HSD) multiple comparison).

Spatial transcript profiles

The spatial distribution of *Asutdes1* and *Asutdes2* in different tissues (head, MTG, midgut, ovary and fat body) of 10-day-old PE females (calling period) was examined to determine if a biased expression pattern might suggest potential functional roles for the enzyme. Despite encoding the same protein, the two *Asutdes* genes exhibited differential expression with *Asutdes1* expressed in midgut, ovary and fat body albeit at relatively low levels (Fig. 3). In contrast, *Asutdes2* was expressed at varying levels in all tissues examined with maximal expression in the fat body and MTG (Fig. 3). The nearly two orders of magnitude lower expression of the *Asutdes1* transcript in the MTG, the site of *A. suturalis* SP biosynthesis and release (Zhang *et al.*, 2016), suggests that the *Asutdes2* gene may play a more important role in *A. suturalis* SP biosynthesis.

The relative expression of *Asutdes* after RNAi-mediated knockdown

To examine the *in vivo* role of *Asutdes* on SP biosynthesis, double-stranded RNA (dsRNA) corresponding to a fragment of the coding sequence shared between the two genes (Fig. 1A) was synthesized and injected into day 0 PE females. The impact of RNAi knockdown was studied at the transcriptional level by quantitative real time PCR (qRT-PCR), using RNA samples from the

MTG and the fat body. These templates were chosen because the MTG is the site of SP synthesis and release (Zhang *et al.*, 2014) and *Asutdes* was most abundant in the fat body, which plays an essential role in energy storage, metabolism and regulation in insects (Arrese & Soulages, 2010). Compared with the control females injected with dsRNA for green fluorescent protein (dsGFP), the dsRNA *Asutdes* (*dsAsutdes*) females had significantly reduced levels of mRNA encoding *Asutdes1*, *Asutdes2* and global *Asutdes* (both *Asutdes1* and *Asutdes2*) in the fat body at 3, 5, 7 and 10 days postinjection (PI) ($P < 0.05$), with transcript reduction ranging from 71–92% (Fig. 4C–E). In the MTG, the relative expression levels of *Asutdes2* and global *Asutdes* were significantly reduced by 45–68% at 3, 7 and 10 days PI ($P < 0.05$), but reduced by 27 and 15%, respectively, at 5 days PI relative to the dsGFP control (Fig. 4A, B). As *Asutdes1* transcript levels in control MTG are barely detectable, its RNAi-mediated silencing was essentially negligible. Gene knockdown in females at 7 days PI showed the best gene silencing efficiency, which coincides well with the reported calling period, 7–10 days PE, of *A. suturalis* females (Zhang *et al.*, 2011). Based on these results, 7 days PI was selected as a suitable period for subsequent RNAi studies.

Knockdown of *Asutdes* clearly suppressed the sexual attractiveness of *A. suturalis* females

Mate-finding behaviour in many insects is largely influenced by pheromones. Consequently, to examine the effect of *Asutdes* silencing on the sexual attractiveness of *A. suturalis* females, both Y-tube olfactometer bioassays and field trapping experiments were performed. For the Y-tube olfactometer bioassay, a total of 174 virgin males (7 days PE) were tested. Given a choice between control females injected with dsGFP and females injected with *dsAsutdes*, males exhibited a statistically significant higher preference for the control females (66.7%, $\chi^2 = 16.67$, $df = 1$, $P < 0.01$; Fig. 5A). The impact of *dsAsutdes* knockdown on the sexual attractiveness of females was also tested in field trapping experiments in both 2015 and 2016. All of the plant bugs that were attracted by the *dsAsutdes*-injected and control females were males, with the number of males attracted significantly reduced in the *dsAsutdes*-injected females relative to the dsGFP-injected and untreated females by 34–53 and 46–59%, respectively ($P < 0.05$; Fig. 5B). The differences between dsGFP-injected and untreated groups were not significant (Fig. 5B). Taken together, knockdown of *Asutdes* had clear suppression effects on the sexual attractiveness of *A. suturalis* females in both laboratory and field test experiments.




Figure 4. Relative mRNA abundance of *Adelphocoris suturalis desaturase* (*Asutdes*) following double-stranded RNA (dsRNA) injection. Virgin females at day 0 posteclosion (PE) were microinjected with either dsRNA for *green fluorescent protein* (*dsGFP*) or *dsAsutdes*, which corresponds to a fragment shared by the two *dsAsutdes* transcripts. After *dsAsutdes* injection, transcript levels in the metathoracic scent gland (MTG) (A, B) and fat body (FB) (C, D and E) were detected at 3, 5, 7 and 10 days postinjection to determine the knockdown efficiency of *Asutdes1* (D), *Asutdes2* (B, E) and both *Asutdes* transcripts, defined as global *des* (A, C). The values are expressed as the means \pm SEM based on three independent biological replicates. The asterisks indicate statistical significance (*, $P < 0.05$; **, $P < 0.01$, Student's *t*-test).

Knockdown of *Asutdes* altered the *A. suturalis* female pheromone titre

Given the reduced female sexual attractiveness following *Asutdes* knockdown, we next examined the effect of *Asutdes* RNAi-mediated silencing on MTG pheromone components. GC-MS analysis of *dsAsutdes*-injected females revealed a notable increase in E4O2H compared with the *dsGFP*-injected controls ($P < 0.05$; Fig. 6C, D). In addition, there was also a slight but nonsignificant increase in HH levels in *dsAsutdes*-injected females ($P = 0.426$; Fig. 6C). The relative ratio between E4O2H and HH in *dsAsutdes*-injected females was 0.90:1 compared to 0.28:1 in the *dsGFP*-injected controls with a significant difference between the two groups ($P < 0.01$; Fig. 6C). Taken together, these results demonstrated that knockdown of *Asutdes* enhances the production of E4O2H and changes the ratio of the *A. suturalis* SP blend.

Discussion

The molecular mechanisms of pheromone biosynthesis have been studied extensively in Lepidoptera (Lassance *et al.*, 2010), Coleoptera (Keeling *et al.*, 2013), Diptera

(Chertemps *et al.*, 2006) and Hymenoptera (Buček *et al.*, 2013). In contrast, little is known about the mechanisms underlying pheromone biosynthesis in Hemiptera. In this study, we isolated and characterized two new *desaturase*-like genes, *Asutdes1* and *Asutdes2*, from *A. suturalis*. RNAi-mediated suppression of *Asutdes* significantly enhanced the production of E4O2H, altered the *A. suturalis* SP blend ratio, and suppressed male attractiveness to *A. suturalis* females. These data strongly support a role for *Asutdes* function in *A. suturalis* SP biosynthesis.

Desaturases are the primary group of proteins responsible for introducing carbon-carbon double bonds at specific positions in fatty acyl substrates. They are found in various eukaryotes including animals, plants, yeast, fungi and many bacteria (Los & Murata, 1998; Sperling *et al.*, 2003). They are important for basic biological processes, including cell signalling, lipid metabolism and regulation of cell membrane fluidity (Hazel & Williams, 1990; Vigh *et al.*, 1993; Pyne & Pyne, 2000; Miyazaki & Ntambi, 2003). In insects, desaturases have been extensively studied based on their role in chemical communication and potential to contribute to speciation. To date, a number of desaturase genes have been shown to be




Figure 5. Effects of RNA interference (RNAi) treatment on the sexual attractiveness of *Adelphocoris suturalis* females. (A) Y-tube olfactometer behavioural response of *A. suturalis* males (7 days posteclosion) to females 7 days postinjection (PI) with either double-stranded RNA (dsRNA) for *A. suturalis* desaturase (*dsAsutdes*) or for green fluorescent protein (*dsGFP*). The number of males that did not make a choice within the 10-min time period is shown as 'not responding'. The total number of tested males was 174. The data were analysed using a chi-square test (***, $P < 0.001$); males that did not respond were not included in the statistical analysis. (B) The number of field-collected males attracted to dsRNA-injected *A. suturalis* females 7–10 days PI. For the RNAi treatment, females were injected with *dsAsutdes*, whereas control females were injected with *dsGFP* or not injected. The values are expressed as the means \pm SEM based on three independent biological replicates. The asterisks indicate statistical significance (*, $P < 0.05$; **, $P < 0.01$, Student's *t*-test). CK, not injected.

involved in pheromone biosynthesis in many insects including moths (Roelofs & Rooney, 2003; Buček *et al.*, 2015) and flies (Knipple *et al.*, 2002; Buček *et al.*, 2013). In these systems, the introduction of carbon–carbon double bonds at specific positions in pheromone precursors to generate pheromone components that vary in chain length, double-bond number, double-bond position and double-bond orientation contributes to the structural diversity in pheromone structures that drive species specificity (Knipple *et al.*, 2002; Roelofs & Rooney, 2003). The two putative desaturase genes *Asutdes1* and *Asutdes2* that we identified appear to be the result of gene duplication rather than alternative splicing as Southern blots exhibited two distinct bands corresponding to each gene. Furthermore, the spatial distribution profiles of the two gene products revealed that *Asutdes2* was most abundant in the MTG and fat body. In contrast, the levels of *Asutdes1* transcripts were low in the

five tissues assayed, in particular the MTG in which expression was lowest (Fig. 3). Based on this differential expression, we hypothesize that *Asutdes2* may play a more important role in *A. suturalis* SP biosynthesis.

Given the structural similarities between moth SPs and those described in *A. suturalis*, we hypothesized that the pheromone biosynthetic pathways might also be similar. We initially identified four genes in *A. suturalis* as candidate pheromone biosynthesis genes based on sequence homology with key enzymes in moth SP biosynthesis (Luo *et al.*, 2014). In this study, we found that silencing *Asutdes* expression potentially suppressed the sexual attractiveness of males to females in both laboratory (Y-tube olfactometer) and field conditions. Further analysis revealed that *Asutdes* knockdown significantly increased levels of E4O2H in the MTG and by doing so altered the *in vivo* ratio of the SP blend (ie E4O2H:HH) by threefold. In insects, the ratios of components in SP blends are highly species specific such that even slight changes have significant effects on male attractiveness (Phelan, 1992; Symonds & Elgar, 2008).

The pheromone biosynthetic pathways of many insects are derived from conventional metabolic pathways (Jurenka, 2004). For example, in moths, the backbones of most pheromone components are generated via fatty acid biosynthesis and then undergo various modifications prior to release (Knipple *et al.*, 2002; Jurenka, 2004; Groot *et al.*, 2016). In *D. melanogaster*, Z9-hexadecenoic is used as a precursor for both ω 7 fatty acid biosynthesis and unsaturated hydrocarbon sex pheromones (Ueyama *et al.*, 2005). Consequently, disruption of one part of the pathway typically results in the accumulation of compounds upstream of the affected site. Furthermore, SP biosynthetic pathways are controlled both by the flux of precursor products through the pathway and enzyme regulation (Lassance *et al.*, 2010; Blaul *et al.*, 2014). We thus speculate that the increase in E4O2H content might be an indication that *Asutdes* functions downstream of E4O2H. In this scenario, *Asutdes* would modulate the rate of substrate movement through the SP biosynthetic pathway by using either E4O2H or a compound downstream of E4O2H as a substrate. The biologically relevant ratio of E4O2H in the MTG would thus be maintained via the controlled flux of substrates, and conversely disruption of *Asutdes* function would impede the flux and trigger an accumulation of E4O2H and the associated disequilibrium of the E4O2H:HH ratio. This proposed function is similar to that of ipsdienol dehydrogenase in *Ips pini*, which oxidizes ipsdienol (a component of the SP in *I. pini*) to ipsdienone, which is then consumed to maintain the biologically active enantiomeric ratio of ipsenol (Rubi *et al.*, 2012). A second potential mechanism for the observed biological effects is that *Asutdes* could function




Figure 6. Effect of *Adelphocoris suturalis desaturase* (*Asutdes*) knockdown on *A. suturalis* female sex pheromone titres. Graphs represent gas chromatogram traces for total ion currents of *n*-hexane extracted metathoracic scent glands from *A. suturalis* females injected with double-stranded RNA for *Asutdes* (*dsAsutdes*; A) or for *green fluorescent protein* (*dsGFP*; B). The internal standard (IS) was *cis*-3-hexenyl isobutyrate. (C) Compared with females injected with *dsGFP*, there was a notable increase in the levels of (E)-4-oxo-2-hexenal (E4O2H) in extracts from *dsAsutdes*-injected females ($P < 0.05$). There was a significant difference in the relative E4O2H : hexyl hexanoate (HH) ratio between *dsAsutdes*-injected females and the *dsGFP* controls. The *y*-axis represents the relative abundance. (D) The total amounts of E4O2H were calculated with the internal standard method in CHEMSTATION. All values are expressed as the means \pm SEM based on six independent biological replicates. A Shapiro–Wilk test was used to test for normality ($P > 0.05$) with significant differences determined using Student's *t*-test (*, $P < 0.05$; **, $P < 0.01$).

in a non-SP pathway that utilizes the same pool of precursors as SP biosynthesis such that *Asutdes* knockdown would result in an accumulation of precursors that would then be shunted exclusively into the SP pathway and converted at a greater rate than normal into E4O2H. Although these hypotheses are potentially verifiable by GC-MS, we did not observe any significant changes in the MTG chemical profile aside from the increase in E4O2H.

Alternatively, silencing *Asutdes* expression might lead to an overexpression of other genes that contribute to E4O2H production. Although the specific involvement of desaturases in this type of regulatory mechanism has not been reported, feedback loops are common. For example, in the migratory brown planthopper, *Nilaparvata lugens*, high *Nilaparvata lugens insulin receptor1* (*NilInR1*) signalling activity inhibits *Nilaparvata lugens forkhead transcription factor Foxo* (*NIFoxo*) activity, leading to long-winged morphs. However, the binding of *NilInR1* to *NilInR2* compromises *NilInR1* signalling activity, which activates *NIFoxo* and results in short-winged morphs (Xu *et al.*, 2015).

Although the precise mechanism underlying the role of *Asutdes* in *A. suturalis* remains to be clarified, our findings provide the first molecular evidence for specific gene function in plant bug SP biosynthesis. Although further exploration of how gene functionality impacts this critical pathway and complex communication systems in

insects is challenging, ongoing advances in molecular techniques are increasing the feasibility. Consequently, future work will focus on exploring the potential molecular processes affected in response to *Asutdes* and elucidating the molecular mechanisms driving the *A. suturalis* SP biosynthetic pathway. Knowledge of these systems can then be applied to further our understanding of similar processes in other hemipteran pests.

Experimental procedures

Insect rearing

A. suturalis were originally collected from a Bt cotton field located in Wuhan (Hubei Province, China) in July 2014. Nymphs and adults were maintained in plastic cages (22.5 \times 15 \times 11 cm) and reared on a 5% sugar solution and green beans, which also served as oviposition substrates (Lu *et al.*, 2008b). All insects were placed in environmental chambers maintained at 75 \pm 5% relative humidity, 26 \pm 2 $^{\circ}$ C temperature and a 16:8 h light : dark (L : D) cycle. The newly emerged adults were separated by sex daily and designated as 0 days PE. All of the experiments performed in this study used virgin females and males.

cDNA cloning and sequence analysis

Total RNA was isolated from *A. suturalis* females 7–10 days PE using a SV total RNA isolation system (Promega, Madison, WI, USA) according to the manufacturer's protocol. RNA integrity was evaluated on 1.5% agarose gels and quantified using a

Table 1. PCR primers used in this study

Primers	Primer sequence	Product size (bp)	PCR efficiency (%)	Standard curve R ²
For cDNA cloning				
<i>Asutdes1</i> -F	CGGGCATCCGAGATTTCAC	1863	n.a.	n.a.
<i>Asutdes1</i> -R	ACAGAGCAAGAACGGTGGTG			
<i>Asutdes2</i> -F	GCGTCTCGCGTTTAAACTTTTC	1780	n.a.	n.a.
<i>Asutdes2</i> -R	ACAGAGCAAGAACGGTGGTG			
For real-time PCR				
Q- <i>Asutdes1</i> -F	TCCGAGATTTACGTTGTCGCA	127	99.5	0.997
Q- <i>Asutdes1</i> -R	AGCAAAATGCTGAAGTGAAGTGCAC			
Q- <i>Asutdes2</i> -F	CTTTTACAGCGCCCTGACCTG	145	94.0	0.997
Q- <i>Asutdes2</i> -R	GCAGGGGCTCCGCTTGTATG			
Q- <i>Asutdes</i> -F	GCTCCCTCTGAGATTGCTGC	130	97.6	0.999
Q- <i>Asutdes</i> -R	TGGTAGCGTTGTGAGGGTGC			
Q- <i>RPS15</i> -F	TGGTTGGCAGCATTGTCGGTG	137	92.6	0.997
Q- <i>RPS15</i> -R	ACCAATACCGGGCCTTCCGT			
Q- <i>EF1γ</i> -F	TTGGCCCTTGCTGCAGAACC	170	91.0	0.998
Q- <i>EF1γ</i> -R	TCTCCGAGCCAGATGGAGTAGTT			
For dsRNA synthesis				
<i>dsAsutdes</i> -F	gogtaatacagactactatagg (T7 promoter) CTATAGCGATGGCCCCAAC	324	n.a.	n.a.
<i>dsAsutdes</i> -R	gogtaatacagactactatagg (T7 promoter) GCAATCTCAGAGGGAGCCTG			
<i>dsGFP</i> -F	gogtaatacagactactatagg (T7 promoter) TGGTCCAATTCTCGTGGAAC	467	n.a.	n.a.
<i>dsGFP</i> -R	gogtaatacagactactatagg (T7 promoter) CTTGAAGTTGACCTTGATGCC			
For <i>Asutdes</i> probe synthesis				
p- <i>Asutdes</i> -F	CTCCCTCTGAGATTGCTGCT	599	n.a.	n.a.
p- <i>Asutdes</i> -R	ATTCTGGCCATGAAGTCGAT			

Asutdes, *Adelphocoris suturalis* desaturase; dsRNA, double-stranded RNA; *EF1 γ* , elongation factor-1 γ ; *GFP*, green fluorescent protein; n.a., not applicable; *RPS15*, ribosomal protein 15.

Nano-Drop 2000 (Thermo Scientific, Wilmington, DE, USA). For first-strand cDNA synthesis, 1 μ g RNA was reverse transcribed using PrimeScriptTM RT Master Mix (perfect real time) (Takara, Kyoto, Japan). Full-length *Asutdes1* and *Asutdes2* were amplified using primers (Table 1) designed based upon *A. suturalis* transcriptome data (Luo *et al.*, 2014), cloned into a pEASY-T1 Simple Cloning Kit (TransGen, Beijing, China) and sequenced. The ExPASy translate tool (<http://web.expasy.org/translate/>) was used to deduce the amino acid sequence of the protein encoded by the *Asutdes* ORF. The full-length *Asutdes* sequence and desaturase sequences from 40 insect species (accession numbers in Table S1) were used to construct a rooted phylogenetic tree with the maximum-likelihood method in MEGA 7.1 (Kumar *et al.*, 2016). Statistical support for the tree was carried out using bootstrap analysis of 1000 replicates. The protein sequence alignments were formatted in DNAMAN 6.0 using the ClustalX colour scheme (Lynnon, San Ramon, CA, USA). The SMART webserver tool (<http://smart.embl.de/>) was used to predict functional domains. The *A. suturalis* cDNA sequences have been deposited with GenBank under the following accession numbers: *Asutdes1*, KX765193; *Asutdes2*, KX765194.

Southern blot

Genomic DNA was extracted from *A. suturalis* females 7–10 days PE using Insect DNA Kit (Omega Bio-tek, Norcross, GA, USA) and a 30 μ g aliquot was digested for 48 h with *HindIII*-HF

and electrophoresed on a 0.8% agarose gel. The separated DNA fragments were blotted onto a Hybond N+ nylon membrane (Millipore, Darmstadt, Germany) and a common region of the two *Asutdes* genes, labelled with digoxigenin, was used as a probe to hybridize with the digested genomic DNA. Southern hybridization was performed using a DIG-High Prime DNA Labeling and Detection Starter Kit II (Roche, Mannheim, Germany), according to the manufacturer's instructions. The primers used to generate the *Asutdes* probe are listed in Table 1.

qRT-PCR

We performed qRT-PCR according to the Minimum Information for publication of Quantitative real time PCR Experiments guidelines (Bustin *et al.*, 2010). All qRT-PCR reactions were performed in a 20 μ l reaction volume using a 20-fold dilution of cDNA with qRT-PCR primers (Table 1) and SYBR[®] Premix ExTaqTM II (Takara) on a Bio-Rad Detection iQ2 System (Bio-Rad, Hercules, CA, USA). PCR conditions consisted of: 95 $^{\circ}$ C for 30 s, followed by 40 cycles of 95 $^{\circ}$ C for 5 s and 60–64 $^{\circ}$ C for 35 s. The qRT-PCR primers were designed using an online tool (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) and the specificity of the PCR products was verified by direct sequencing. PCR efficiencies (Table 1) were determined using standard curves generated from serially diluted cDNAs and the specificity of the reaction confirmed by melt curve analysis (Liu *et al.*, 2016). Each reaction was performed in triplicate using at least three biological replicates per individual treatment. The qRT-

PCR data were analysed via the $2^{-\Delta\Delta Ct}$ method (Schmittgen & Livak, 2008).

Spatial expression pattern analyses

To investigate the spatial expression profile of the *Asutdes* transcripts, total RNAs were collected from the head, MTG, midgut, ovary and fat body of females 10 days PE (calling period; Zhang *et al.*, 2011). cDNA synthesis and mRNA quantification were performed as described above. *Ribosomal protein S15* was used as the reference gene for gene expression normalization (unpubl. data). The values are expressed as the means \pm SEM. The data were analysed using one-way analysis of variance followed by Tukey's Honestly Significant Difference (HSD) multiple comparison test with SPSS 18.0 (SPSS Inc., Chicago, IL, USA). Statistical differences are indicated by different letters in the figures.

RNAi in *A. suturalis*

dsRNAs corresponding to a 324-bp fragment of the coding region shared by the *Asutdes1* and *Asutdes2* genes were synthesized as described (Liu *et al.*, 2016). Similarly, *dsGFP* was synthesized and used as the control. dsRNA integrity was evaluated on 1.5% agarose gels and quantified on a Nano-Drop 2000 (Thermo Scientific). *A. suturalis* females at 0 day PE were microinjected with 2 μ g dsRNA per 200 nl. Total RNAs from the MTG and fat body were extracted 3, 5, 7 and 10 days PI to determine the RNAi efficiency via qRT-PCR. *Elongation factor-1 γ* was used as the reference gene for gene expression normalization (unpubl. data). Significant differences were determined using Student's *t*-test (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$). Based on these results, females 7 days PI were used in all subsequent bioassays.

Y-tube olfactometer bioassay

To evaluate the biological effect of *Asutdes* knockdown, the sexual attractiveness of *A. suturalis* females was assayed in a Y-tube olfactometer system. Mature males (7 days PE) were assayed for behavioural responses to odours from two experimental groups that consisted of five females injected 7 days prior with either *dsAsutdes* or *dsGFP*. The olfactometer set-up consisted of a Y-shaped Pyrex tube (Wuhan, China) (internal diameter 30 mm), with a 220 mm central stem and two 160 mm side arms set at a 90° angle. Air was moved through the system using silicon tubing attached to an air pump (Beijing Municipal Institute of Labor Protection, Beijing, China). An activated charcoal filter was used to purify the air, which was humidified by bubbling through distilled water prior to being split with a glass Y-connector. The air flow rate of each branch was set to 150 ml/min with a flow meter (Huangming, Zhejiang, China) and then passed through two modified jars (100 ml volume capacity), each containing the odour source, before being passed through the arms of the Y-tube. In each test, a single mature male was released at the entrance of the central stem of the Y-tube. All insects were reared under a photoperiod of 16:8 h (L : D) and maintained in the dark for at least 1–2 h after lights out. In the olfactometer set-up, movement of the tested males beyond the decision line (2 cm up either side arm) for a

minimum of 20 s was recorded as a response, whereas males that failed to move within 10 min were recorded as 'no response' (Bodino *et al.*, 2016). Each male was tested only once to avoid any conditioned behaviour and the test odour sources were changed after every 10 tested males. To compensate for spatial asymmetry in the set-up, we switched the odour sources after testing four males. The Y-tube was cleaned thoroughly with hexane and baked before each individual test. All trials were performed at the same time (from 8 pm to 2 am). Olfactory bioassays were conducted in the dark (with a red light) at 26 ± 1 °C and $45 \pm 10\%$ relative humidity (Zhang *et al.*, 2011). The data were analysed using a chi-square test (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$); males that did not respond were not included in the statistical analysis.

Field trapping experiments

Field trapping experiments were carried out in transgenic cotton fields in Wuhan (Hubei Province, China) during August of both 2015 and 2016. The number of adults assayed per 100 plants per year was 21 and 19, respectively. An equilateral-triangular-prism trap (16 cm on each side and 19 cm in height) was used for this study. Adhesive glue was painted on the inner surface of the traps to catch insects that entered. Three females injected with *dsAsutdes* (7 days PI) were placed in a small nylon cage (3.5 \times 3.5 \times 3.5 cm) and used as lures, with one young bud of cotton inside for female physical activity. The lures were hung in the middle of the traps, 2 cm above the base. Traps were placed randomly in the cotton field, 1.5 m above the ground and a minimum of 15 m apart (Zhang *et al.*, 2011). Females injected with *dsGFP* or that were untreated were used as controls. All traps were placed in the cotton field at 5 pm and removed at 7 am the next morning, at which time the number of trapped adults was recorded. Each female was tested only once. The field trapping experiments were performed for at least 1 week with three biological replicates performed for each individual treatment in the field. A Student's *t*-test was used to test for significant differences (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Pheromone extraction and GC-MS analysis

Extracts were prepared by placing 10 MTGs from dsRNA-injected females (7 days PI) in 100 μ l *n*-hexane (Aladdin, Shanghai, China purity > 98%) for 30 min at room temperature with an internal standard, which consisted of 50 ng/ml cis-3-hexenyl isobutyrate (Sigma, St Louis, MO, USA, purity > 98%). MTG extracts were then stored at -20 °C until needed. GC-MS analyses were performed on an Agilent 7890B GC (Agilent Technologies, Palo Alto, CA, USA) coupled to a mass detector (Agilent 5977C) equipped with a HP-5 capillary column (30 m \times 0.25 mm). Six independent biological replicates were performed for each treatment. GC-MS conditions consisted of: He as the carrier gas at 1 ml/min; 1 μ l splitless injection; 220 °C injector temperature; thermal gradient of 50 °C for 1 min then 50 to 250 °C at 10 °C/min with termination at 280 °C for 2 min. Data were analysed using CHEMSTATION software (Agilent Technologies). HH was purchased from Sigma (purity > 98%) and E4O2H (purity > 95%) was donated from Tao Zhang (State Key Laboratory for Biology of Plant Diseases and

Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China). Chromatograms shown are representative of the extract analyses. A Shapiro–Wilk test was used to test for normality ($P > 0.05$) and a Student's *t*-test was used to test for significant differences (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

Acknowledgements

This work was funded by grants from the Ministry of Agriculture of China (grant no. 2016ZX08011002–006) and State Key Laboratory for Biology of Plant Diseases and Insect Pests (grant no. SKLOF201415). Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the United States Department of Agriculture. USDA is an equal opportunity provider and employer.

References

- Albre, J., Liénard, M.A., Sirey, T.M., Schmidt, S., Tooman, L.K., Carraher, C. *et al.* (2012) Sex pheromone evolution is associated with differential regulation of the same desaturase gene in two genera of leafroller moths. *PLoS Genet* **8**: e1002489.
- Antony, B., Ding, B.J., Moto, K., Aldosari, S.A. and Aldawood, A.S. (2016) Two fatty acyl reductases involved in moth pheromone biosynthesis. *Sci Rep* **6**: 29927.
- Arrese, E.L. and Soulages, J.L. (2010) Insect fat body: energy, metabolism, and regulation. *Annu Rev Entomol* **55**: 207–225.
- Blaul, B., Steinbauer, R., Merkl, P., Merkl, R., Tschochner, H. and Ruther, J. (2014) Oleic acid is a precursor of linoleic acid and the male sex pheromone in *Nasonia vitripennis*. *Insect Biochem Mol Biol* **51**: 33–40.
- Bodino, N., Ferracini, C. and Tavella, L. (2016) Is host selection influenced by natal and adult experience in the parasitoid *Necremnus tutae* (Hymenoptera: Eulophidae)? *Anim Behav* **112**: 221–228.
- Buček, A., Vogel, H., Matoušková, P., Prchalová, D., Žáček, P., Vrkoslav, V. *et al.* (2013) The role of desaturases in the biosynthesis of marking pheromones in bumblebee males. *Insect Biochem Mol Biol* **43**: 724–731.
- Buček, A., Matoušková, P., Vogel, H., Šebesta, P., Jahn, U., Weißflog, J. *et al.* (2015) Evolution of moth sex pheromone composition by a single amino acid substitution in a fatty acid desaturase. *Proc Natl Acad Sci USA* **112**: 12586–12591.
- Bustin, S.A., Beaulieu, J.F., Huggett, J., Jaggi, R., Kibenge, F.S.B., Olsvik, P.A. *et al.* (2010) MIQE précis: practical implementation of minimum standard guidelines for fluorescence-based quantitative real-time PCR experiments. *BMC Mol Biol* **11**: 1–5.
- Chertemps, T., Duportets, L., Labeur, C., Ueyama, M. and Wicker-Thomas, C. (2006) A female-specific desaturase gene responsible for diene hydrocarbon biosynthesis and courtship behaviour in *Drosophila melanogaster*. *Insect Mol Biol* **15**: 465–473.
- Chertemps, T., Duportets, L., Labeur, C., Ueda, R., Takahashi, K., Saigo, K. *et al.* (2007) A female-biased expressed elongase involved in long-chain hydrocarbon biosynthesis and courtship behavior in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **104**: 4273–4278.
- Chung, H., Loehlin, D.W., Dufour, H.D., Vaccarro, K., Millar, J.G. and Carroll, S.B. (2014) A single gene affects both ecological divergence and mate choice in *Drosophila*. *Science* **343**: 1148–1151.
- Coyne, J.A. and Charlesworth, B. (1997) Genetics of a pheromonal difference affecting sexual isolation between *Drosophila mauritiana* and *D. sechellia*. *Genetics* **145**: 1015–1030.
- Dallerac, R., Labeur, C., Jallon, J.M., Knipple, D.C., Roelofs, W.L. and Wicker-Thomas, C. (2000) A $\Delta 9$ desaturase gene with a different substrate specificity is responsible for the cuticular diene hydrocarbon polymorphism in *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **97**: 9449–9454.
- EI-Sayed, A. M. (2016) *The Pherobase: Database of Pheromones and Semiochemicals*. <http://www.pherobase.com>. Accessed on 14 December 2016.
- Fujii, T., Ito, K., Tatematsu, M., Shimada, T., Katsuma, S. and Ishikawa, Y. (2011) Sex pheromone desaturase functioning in a primitive *Ostrinia* moth is cryptically conserved in congeners' genomes. *Proc Natl Acad Sci USA* **108**: 7102–7106.
- González-Caballero, N., Rodríguez-Vega, A., Dias-Lopes, G., Valenzuela, J.G., Ribeiro, J.M.C., Carvalho, P.C. *et al.* (2014) Expression of the mevalonate pathway enzymes in the *Lutzomyia longipalpis* (Diptera: Psychodidae) sex pheromone gland demonstrated by an integrated proteomic approach. *J Proteomics* **96**: 117–132.
- Groot, A.T., Schöfl, G., Inglis, O., Donnerhacke, S., Classen, A., Schmalz, A. *et al.* (2014) Within-population variability in a moth sex pheromone blend: genetic basis and behavioural consequences. *Proc R Soc B* **281**: 20133054.
- Groot, A.T., Dekker, T. and Heckel, D.G. (2016) The genetic basis of pheromone evolution in moths. *Annu Rev Entomol* **61**: 99–117.
- Hagström, A.K., Albre, J., Tooman, L.K., Thirmawithana, A.H., Corcoran, J., Lofstedt, C. *et al.* (2014) A novel fatty acyl desaturase from the pheromone glands of *Ctenopseustis obliquana* and *C. herana* with specific Z5-desaturase activity on myristic acid. *J Chem Ecol* **40**: 63–70.
- Hao, G., O'Connor, M., Liu, W. and Roelofs, W.L. (2002) Characterization of Z/E11- and Z9-desaturases from the obliquebanded leafroller moth, *Choristoneura rosaceana*. *J Insect Sci* **2**: 26.
- Haritos, V.S., Horne, I., Damcevski, K., Glover, K. and Gibb, N. (2014) Unexpected functional diversity in the fatty acid desaturases of the flour beetle *Tribolium castaneum* and identification of key residues determining activity. *Insect Biochem Mol Biol* **51**: 62–70.
- Hazel, J.R. and Williams, E.E. (1990) The role of alterations in membrane lipid composition in enabling physiological adaptation of organisms to their physical environment. *Prog Lipid Res* **29**: 167–227.
- Johansson, B.G. and Jones, T.M. (2007) The role of chemical communication in mate choice. *Biol Rev Camb Philos Soc* **82**: 265–289.
- Jurenka, R. (2004) Insect pheromone biosynthesis. *Top Curr Chem* **239**: 97–132.
- Jurenka, R.A., Haynes, K.F., Adlof, R.O., Bengtsson, M. and Roelofs, W.L. (1994) Sex pheromone component ratio in the cabbage looper moth altered by a mutation affecting the fatty acid chain-shortening reactions in the pheromone biosynthetic pathway. *Insect Biochem Mol Biol* **24**: 373–381.

- Keeling, C.I., Chiu, C.C., Aw, T., Li, M., Henderson, H., Tittiger, C. *et al.* (2013) Frontalin pheromone biosynthesis in the mountain pine beetle, *Dendroctonus ponderosae*, and the role of isoprenyl diphosphate synthases. *Proc Natl Acad Sci USA* **110**: 18838–18843.
- Knipple, D.C., Rosenfield, C.L., Nielsen, R., You, K.M. and Jeong, S.E. (2002) Evolution of the integral membrane desaturase gene family in moths and flies. *Genetics* **162**: 1737–1752.
- Kumar, S., Stecher, G. and Tamura, K. (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* **33**: 1870–1874.
- Lassance, J.M., Groot, A.T., Lienard, M.A., Antony, B., Borgwardt, C., Andersson, F. *et al.* (2010) Allelic variation in a fatty-acyl reductase gene causes divergence in moth sex pheromones. *Nature* **466**: 486–489.
- Lassance, J.M., Liénard, M.A., Antony, B., Qian, S., Fujii, T., Tabata, J. *et al.* (2013) Functional consequences of sequence variation in the pheromone biosynthetic gene pgFAR for *Ostrinia* moths. *Proc Natl Acad Sci USA* **110**: 3967–3972.
- Li, G.P., Feng, H.Q., Chen, P.Y., Wu, S.Y., Liu, B. and Qiu, F. (2010) Effects of transgenic Bt cotton on the population density, oviposition behavior, development, and reproduction of a nontarget pest, *Adelphocoris suturalis* (Hemiptera: Miridae). *Environ Entomol* **39**: 1378–1387.
- Liu, W., Ma, P.W.K., Marsella-Herrick, P., Rosenfield, C.L., Knipple, D.C. and Roelofs, W. (1999) Cloning and functional expression of a cDNA encoding a metabolic acyl-CoA $\Delta 9$ -desaturase of the cabbage looper moth, *Trichoplusia ni*. *Insect Biochem Mol Biol* **29**: 435–443.
- Liu, W., Li, Y., Zhu, L., Zhu, F., Lei, C.L. and Wang, X.P. (2016) Juvenile hormone facilitates the antagonism between adult reproduction and diapause through the methoprene-tolerant gene in the female *Colaphellus bowringi*. *Insect Biochem Mol Biol* **74**: 50–60.
- Los, D.A. and Murata, N. (1998) Structure and expression of fatty acid desaturases. *Biochim Biophys Acta* **1394**: 3–15.
- Lu, Y.H., Qiu, F., Feng, H.Q., Li, H.B., Yang, Z.C., Wyckhuys, K.A.G. *et al.* (2008a) Species composition and seasonal abundance of pestiferous plant bugs (Hemiptera: Miridae) on Bt cotton in China. *Crop Prot* **27**: 465–472.
- Lu, Y.H., Wu, K.M., Cai, X.M. and Liu, Y.Q. (2008b) A rearing method for mirids using the green bean, *Phaseolus vulgaris* in the laboratory. *Acta Phytophysiol Sinica* **35**: 251–269.
- Lu, Y.H., Wu, K.M., Jiang, Y.Y., Xia, B., Li, P., Feng, H.Q. *et al.* (2010) Mirid bug outbreaks in multiple crops correlated with wide-scale adoption of Bt cotton in China. *Science* **328**: 1151–1154.
- Luo, J., Liu, X.Y., Liu, L., Zhang, P.Y., Chen, L.J., Gao, Q. *et al.* (2014) De novo analysis of the *Adelphocoris suturalis* Jakovlev metathoracic scent glands transcriptome and expression patterns of pheromone biosynthesis-related genes. *Gene* **551**: 271–278.
- Miyazaki, M. and Ntambi, J.M. (2003) Role of stearoyl-coenzyme A desaturase in lipid metabolism. *Prostaglandins Leukot Essent Fatty Acids* **68**: 113–121.
- Moto, K., Yoshiga, T., Yamamoto, M., Takahashi, S., Okano, K., Ando, T. *et al.* (2003) Pheromone gland-specific fatty-acyl reductase of the silkworm, *Bombyx mori*. *Proc Natl Acad Sci USA* **100**: 9156–9161.
- Moto, K., Suzuki, M.G., Hull, J.J., Kurata, R., Takahashi, S., Yamamoto, M. *et al.* (2004) Involvement of a bifunctional fatty-acyl desaturase in the biosynthesis of the silkworm, *Bombyx mori*, sex pheromone. *Proc Natl Acad Sci USA* **101**: 8631–8636.
- Ng, W.C., Chin, J.S.R., Tan, K.J. and Yew, J.Y. (2015) The fatty acid elongase bond is essential for *Drosophila* sex pheromone synthesis and male fertility. *Nat Commun* **6**: 8263.
- Phelan, P.L. (ed.) (1992) *Insect Chemical Ecology: Evolution of Sex Pheromones and the Role of Asymmetric Tracking*. Chapman & Hall, New York.
- Pyne, S. and Pyne, N.J. (2000) Sphingosine 1-phosphate signaling in mammalian cells. *Biochem J* **349**: 385.
- Roelofs, W.L. and Rooney, A.P. (2003) Molecular genetics and evolution of pheromone biosynthesis in Lepidoptera. *Proc Natl Acad Sci USA* **100**: 9179–9184.
- Roelofs, W.L., Liu, W., Hao, G., Jiao, H., Rooney, A.P. and Linn, C.E. (2002) Evolution of moth sex pheromones via ancestral genes. *Proc Natl Acad Sci USA* **99**: 13621–13626.
- Rosenfield, C.L., You, K.M., Marsella-Herrick, P., Roelofs, W.L. and Knipple, D.C. (2001) Structural and functional conservation and divergence among acyl-CoA desaturases of two noctuid species, the corn earworm, *Helicoverpa zea*, and the cabbage looper, *Trichoplusia ni*. *Insect Biochem Mol Biol* **31**: 949–964.
- Rubi, F.T., Welch, W.H., Blomquist, G.J. and Tittiger, C. (2012) Ipsdienol dehydrogenase (IDOLDH): a novel oxidoreductase important for *Ips pini* pheromone production. *Insect Biochem Mol Biol* **42**: 81–90.
- Sandstrom, P., Welch, W.H., Blomquist, G.J. and Tittiger, C. (2006) Functional expression of a bark beetle cytochrome P450 that hydroxylates myrcene to ipsdienol. *Insect Biochem Mol Biol* **36**: 835–845.
- Schmittgen, T.D. and Livak, K.J. (2008) Analyzing real-time PCR data by the comparative CT method. *Nat Protoc* **3**: 1101–1108.
- Shanklin, J., Whittle, E. and Fox, B.G. (1994) Eight histidine residues are catalytically essential in a membrane-associated iron enzyme, stearoyl-CoA desaturase, and are conserved in alkane hydroxylase and xylene monooxygenase. *Biochemistry* **33**: 12787–12749.
- Song, M., Delaplain, P., Nguyen, T.T., Liu, X., Wickenberg, L., Jeffrey, C. *et al.* (2014) exo-Brevicommin biosynthetic pathway enzymes from the Mountain Pine Beetle, *Dendroctonus ponderosae*. *Insect Biochem Mol Biol* **53**: 73–80.
- Sperling, P., Ternes, P., Zank, T.K. and Heinz, E. (2003) The evolution of desaturases. *Prostaglandins Leukot Essent Fatty Acids* **68**: 73–95.
- Symonds, M.R.E. and Elgar, M.A. (2008) The evolution of pheromone diversity. *Trends Ecol Evol* **23**: 220–228.
- Tabata, J. and Ishikawa, Y. (2005) Genetic basis to divergence of sex pheromones in two closely related moths, *Ostrinia scapularis* and *O. zealis*. *J Chem Ecol* **31**: 1111–1124.
- Tillman, J.A., Seybold, S.J., Jurenka, R.A. and Blomquist, G.J. (1999) Insect pheromones—an overview of biosynthesis and endocrine regulation. *Insect Biochem Mol Biol* **29**: 481–514.
- Ueyama, M., Chertemps, T., Labeur, C. and Wicker-Thomas, C. (2005) Mutations in the desat1 gene reduces the production of courtship stimulatory pheromones through a marked effect on fatty acids in *Drosophila melanogaster*. *Insect Biochem Mol Biol* **35**: 911–920.
- Vigh, L., Los, D.A., Horváth, I. and Murata, N. (1993) The primary signal in the biological perception of temperature: Pd-

- catalyzed hydrogenation of membrane lipids stimulated the expression of the desA gene in *Synechocystis* PCC6803. *Proc Natl Acad Sci USA* **90**: 9090–9094.
- Wang, H.L., Liénard, M.A., Zhao, C.H., Wang, C.Z. and Löfstedt, C. (2010) Neofunctionalization in an ancestral insect desaturase lineage led to rare $\Delta 6$ pheromone signals in the Chinese tussah silkworm. *Insect Biochem Mol Biol* **40**: 742–751.
- Xu, H.J., Xue, J., Lu, B., Zhang, X.C., Zhuo, J.C., He, S.F. *et al.* (2015) Two insulin receptors determine alternative wing morphs in planthoppers. *Nature* **519**: 464–467.
- Xue, B., Rooney, A.P., Kajikawa, M., Okada, N. and Roelofs, W.L. (2007) Novel sex pheromone desaturases in the genomes of corn borers generated through gene duplication and retroposon fusion. *Proc Natl Acad Sci USA* **104**: 4467–4472.
- Zhang, Z.L., Luo, J., Lu, C., Zhao, B., Meng, J., Chen, L.Z. *et al.* (2011) Evidence of female-produced sex pheromone of *Adelphocoris suturalis* (Hemiptera: Miridae): effect of age and time of day. *J Econ Entomol* **104**: 1189–1194.
- Zhang, Z., Luo, J., Wang, Y., Chen, L., Chen, L. and Lei, C. (2014) Morphology and chemical analysis of the metathoracic scent glands system in *Adelphocoris suturalis* (Hemiptera: Miridae). *J Insect Sci* **14**: 293.
- Zhang, Z., Zhang, T., Zhang, A., Luo, J., Chen, L., Wang, M. *et al.* (2016) Identification and field verification of sex pheromone from the mirid bug, *Adelphocoris suturalis*. *Chemoecology* **25**: 25–31.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Table S1. List of species names and corresponding published accession numbers for sequences used in the desaturase phylogeny in Fig. 2B.